

Technical Appendix: Heat Kernel, Fermions, and the Sign of Induced Gravity

(Sign conventions fixed; Laplacian/Lichnerowicz hinge made explicit)

Lluis Eriksson
Independent Researcher
lluiseriksson@gmail.com

December 2025

Scope note. This appendix is a technical support piece for the induced-sign bookkeeping used in the broader “resource boundary / cut” program; see, e.g., [1, 8, 10, 3]. It does not attempt a UV completion nor a physical determination of G_{phys} .

Abstract

We derive the local contribution proportional to the scalar curvature R in the Euclidean one-loop effective action obtained by integrating out matter fields on a curved background. Using a Schwinger cutoff $\varepsilon = \Lambda^{-2}$ and the Seeley–DeWitt coefficient a_1 , we extract the quadratically divergent term multiplying $\int d^4x \sqrt{g} R$. We fix a single Euclidean convention for the Einstein–Hilbert action, state an explicit Laplacian convention, and write the Lichnerowicz/Weitzenböck identity in a sign-robust form so that the fermionic contribution is unambiguous. We provide a unified bookkeeping coefficient $A_1^{(\text{eff})}$ such that

$$W_R^{\text{total}} = -\frac{A_1^{(\text{eff})}}{32\pi^2} \Lambda^2 \int d^4x \sqrt{g} R,$$

and hence an induced Newton constant G_{ind} via comparison with the Euclidean Einstein–Hilbert action. We also include the minimal gauge+ghost package in background Feynman gauge, a species table, and a short reproducible Python snippet.

Keywords: heat kernel, Seeley–DeWitt coefficients, induced gravity, Euclidean QFT in curved space, fermions, ghosts, gauge fixing, Newton constant.

1 Conventions and notation

1.1 Lorentzian geometry and curvature sign

We adopt Lorentzian signature $(-, +, +, +)$ and the Wald/MTW-type curvature convention

$$R^a{}_{bcd} = \partial_c \Gamma_{bd}^a - \partial_d \Gamma_{bc}^a + \Gamma_{ce}^a \Gamma_{bd}^e - \Gamma_{de}^a \Gamma_{bc}^e, \quad R_{bd} = R^a{}_{bad}, \quad R = g^{bd} R_{bd}.$$

1.2 Euclidean formulation and Einstein–Hilbert action

Heat-kernel calculations are performed in $D = 4$ Euclidean signature. We fix the Euclidean Einstein–Hilbert action as

$$S_E^{\text{EH}} = -\frac{1}{16\pi G} \int d^4x \sqrt{g} R,$$

obtained by Wick rotation from the standard Lorentzian action

$$S_L = \frac{1}{16\pi G} \int d^4x \sqrt{-g} R,$$

i.e. $S_E = -iS_L$ when phases from topological/eta-invariants are irrelevant for the local divergent coefficient we extract.

1.3 Laplacian convention, Laplace-type operators, and Schwinger cutoff

Define the covariant Laplacian as

$$\nabla^2 \equiv g^{\mu\nu} \nabla_\mu \nabla_\nu,$$

so that in flat Euclidean space $\nabla^2 = \partial^2$ and $\partial^2 e^{ik \cdot x} = -k^2 e^{ik \cdot x}$. Hence the principal operator $-\nabla^2$ is positive.

A Laplace-type operator is taken as

$$P = -\nabla^2 + X(x),$$

where $X(x)$ is a local endomorphism (a scalar function for scalar fields).

We use Schwinger proper-time regularization with cutoff $\varepsilon = \Lambda^{-2}$:

$$\ln \det P = - \int_\varepsilon^\infty \frac{ds}{s} \text{Tr} e^{-sP}.$$

1.4 Fermionic hinge: commutator and Lichnerowicz/Weitzenböck identity

For spinors we take $D_E \equiv \gamma^\mu \nabla_\mu$ with Euclidean gamma matrices $(\gamma^\mu)^\dagger = \gamma^\mu$ and Dirac operator anti-Hermitian $(D_E)^\dagger = -D_E$. The covariant derivative on spinors includes the spin connection.

We fix the curvature term via

$$[\nabla_\mu, \nabla_\nu] \psi = \frac{1}{4} R_{\mu\nu\rho\sigma} \gamma^{\rho\sigma} \psi, \quad \gamma^{\rho\sigma} \equiv \frac{1}{2} [\gamma^\rho, \gamma^\sigma].$$

With the conventions above, the Lichnerowicz/Weitzenböck identity can be written in the sign-robust form

$$-D_E^2 = -\nabla^2 + \frac{1}{4} R.$$

See, e.g., [12, 13].

Remark 1 (Connection Laplacian). *In the fermionic sector, $\nabla^2 = g^{\mu\nu} \nabla_\mu \nabla_\nu$ is the connection Laplacian acting on spinors, not the de Rham Laplacian on differential forms.*

2 Heat kernel expansion and the coefficient a_1

Let $K(s; x, y) = \langle x | e^{-sP} | y \rangle$ be the heat kernel. The trace has the asymptotic expansion as $s \rightarrow 0^+$:

$$\mathrm{Tr} e^{-sP} \sim \frac{1}{(4\pi s)^2} \int d^4x \sqrt{g} (a_0(x) + a_1(x)s + a_2(x)s^2 + \dots).$$

For a scalar Laplace-type operator $P = -\nabla^2 + X$ the first coefficients are

$$a_0(x) = \mathrm{tr} \mathbf{1}, \quad a_1(x) = \mathrm{tr} \left(\frac{1}{6} R - X \right).$$

For a real scalar with $X = m^2 + \xi R$:

$$a_1(x) = \left(\frac{1}{6} - \xi \right) R - m^2, \quad a_{1,R}(x) = \tilde{a}_1 R, \quad \tilde{a}_1 = \frac{1}{6} - \xi.$$

The term $-m^2$ contributes to $\int \sqrt{g}$ (cosmological constant renormalization) and not to $\int \sqrt{g} R$.

3 Scalars: effective action and extraction of $\int \sqrt{g} R$

For a real bosonic scalar,

$$W_{\mathrm{scalar}} = \frac{1}{2} \ln \det P = -\frac{1}{2} \int_{\varepsilon}^{\infty} \frac{ds}{s} \mathrm{Tr} e^{-sP}.$$

Keeping only the piece proportional to R :

$$W_R^{\mathrm{scalar}} = -\frac{1}{2} \frac{1}{(4\pi)^2} \int d^4x \sqrt{g} a_{1,R}(x) \int_{\varepsilon}^{\infty} ds s^{-2}.$$

Since $\int_{\varepsilon}^{\infty} ds s^{-2} = 1/\varepsilon = \Lambda^2$, we obtain

$$W_R^{\mathrm{scalar}} = -\frac{\tilde{a}_1}{32\pi^2} \Lambda^2 \int d^4x \sqrt{g} R.$$

4 Dirac fermions: reduction to Laplace type and contribution to $\int \sqrt{g} R$

For a Euclidean Dirac fermion with action

$$S_F = \int d^4x \sqrt{g} \bar{\psi} (D_E + m) \psi,$$

the functional integral yields

$$W_{\mathrm{ferm}} = -\ln \det(D_E + m).$$

4.1 Determinant reduction

Using the adjoint product,

$$\ln \det(D_E + m) = \frac{1}{2} \ln \det \left((D_E + m)^\dagger (D_E + m) \right) + i\Theta,$$

where the phase $i\Theta$ does not affect the local divergent coefficient multiplying R . Define the positive operator

$$P_D \equiv (D_E + m)^\dagger (D_E + m).$$

Since $(D_E)^\dagger = -D_E$ and m is constant,

$$P_D = (-D_E + m)(D_E + m) = -D_E^2 + m^2.$$

With Lichnerowicz,

$$P_D = -\nabla^2 + m^2 + \frac{1}{4} R.$$

4.2 The a_1 coefficient and spin trace

Here $X_D = m^2 + \frac{1}{4}R$, so

$$a_1^{(P_D)}(x) = \frac{1}{6}R - X_D = -\frac{1}{12}R - m^2.$$

Tracing over Dirac spin indices in $D = 4$ gives $\text{tr } \mathbf{1} = 4$, hence

$$\text{tr } a_{1,R}^{(P_D)}(x) = 4 \cdot \left(-\frac{1}{12}R\right) = -\frac{1}{3}R.$$

4.3 Schwinger insertion

Because $W_{\text{ferm}} = -\frac{1}{2} \ln \det P_D$ and $\ln \det P_D = -\int ds/s \text{Tr } e^{-sP_D}$,

$$W_{\text{ferm}} = +\frac{1}{2} \int_{\epsilon}^{\infty} \frac{ds}{s} \text{Tr } e^{-sP_D}.$$

Thus the R -term is

$$W_R^{\text{ferm}} = \frac{1}{2} \frac{1}{(4\pi)^2} \int d^4x \sqrt{g} \text{tr } a_{1,R}^{(P_D)}(x) \int_{\epsilon}^{\infty} ds s^{-2} = -\frac{1}{96\pi^2} \Lambda^2 \int d^4x \sqrt{g} R.$$

Remark 2 (Phases and zero modes). *The phase $i\Theta$ and zero modes matter in chiral/topological settings (eta-invariants, anomalies). For the present purpose (local divergent coefficient multiplying R for a massive Dirac field in a smooth background), they do not modify the $\Lambda^2 \int \sqrt{g} R$ term.*

5 Gauge fields and ghosts (spin-1): minimal package in Feynman gauge

For a gauge field (per generator) in background Feynman gauge ($\alpha = 1$), one may take the 1-form operator

$$(P_1)_{\mu}{}^{\nu} = -\delta_{\mu}^{\nu} \nabla^2 + R_{\mu}{}^{\nu},$$

and the Faddeev–Popov ghost operator (complex Grassmann scalar)

$$P_{\text{gh}} = -\nabla^2.$$

The gauge-fixed one-loop contribution has the standard structure

$$W_{\text{gauge}} = \frac{1}{2} \ln \det P_1 - \ln \det P_{\text{gh}}.$$

5.1 Short derivation of $A_1^{(\text{eff})}(\text{vector} + \text{ghosts}) = -\frac{2}{3}$

For a Laplace-type operator on 1-forms, $P_1 = -\nabla^2 + E$ with $E_{\mu}{}^{\nu} = R_{\mu}{}^{\nu}$. Then

$$\text{tr } a_{1,R}^{(1\text{-form})} = \left(\frac{1}{6} \text{tr } \mathbf{1} - \frac{\text{tr } E}{R}\right) R = \left(\frac{4}{6} - 1\right) R = -\frac{1}{3}R.$$

The ghost is a complex Grassmann scalar, so it contributes with the opposite sign of a complex bosonic scalar. Since a complex bosonic scalar gives $\text{tr } a_{1,R} = 2 \cdot (\frac{1}{6}R) = \frac{1}{3}R$, the ghost gives

$$\text{tr } a_{1,R}^{(\text{gh})} = -\frac{1}{3}R.$$

In the unified convention defined below, this yields (per generator)

$$A_1^{(\text{eff})}(\text{vector} + \text{ghosts}) = -\frac{2}{3}.$$

See [12, 15] for detailed heat-kernel treatments in gauge theories.

6 Unified convention and induced Newton constant

Define the unified coefficient $A_1^{(\text{eff})}$ by

$$W_R^{\text{total}} = -\frac{A_1^{(\text{eff})}}{32\pi^2} \Lambda^2 \int d^4x \sqrt{g} R.$$

Comparing with the Euclidean Einstein–Hilbert term

$$S_E^{\text{EH}} = -\frac{1}{16\pi G} \int d^4x \sqrt{g} R,$$

and identifying $S_{E,\text{induced}}^{\text{EH}} = W_R^{\text{total}}$ gives

$$-\frac{1}{16\pi G_{\text{ind}}} = -\frac{A_1^{(\text{eff})}}{32\pi^2} \Lambda^2,$$

hence

$$\boxed{\frac{1}{16\pi G_{\text{ind}}} = \frac{A_1^{(\text{eff})}}{32\pi^2} \Lambda^2 \quad \Rightarrow \quad G_{\text{ind}} = \frac{2\pi}{A_1^{(\text{eff})}} \frac{1}{\Lambda^2}}.$$

With this convention, $G_{\text{ind}} > 0$ if and only if $A_1^{(\text{eff})} > 0$.

7 Species table (unified convention)

Species	$A_1^{(\text{eff})}$	Comment
Real scalar (general ξ)	$\frac{1}{6} - \xi$	$X = m^2 + \xi R$
Real scalar (minimal)	$+\frac{1}{6}$	$\xi = 0$
Complex scalar	$2(\frac{1}{6} - \xi)$	two real d.o.f.
Dirac fermion (4 comp.)	$+\frac{1}{3}$	derived above
Weyl fermion (2 comp.)	$+\frac{1}{6}$	half of Dirac
Majorana fermion	$+\frac{1}{6}$	half of Dirac
Gauge vector + ghosts	$-\frac{2}{3}$	per generator, Feynman gauge

8 Examples and a reproducible Python snippet

8.1 Sign examples

- Minimal real scalar: $A_1^{(\text{eff})} = \frac{1}{6} > 0$ implies $G_{\text{ind}} > 0$.
- Real scalar with $\xi = \frac{1}{4}$: $A_1^{(\text{eff})} = \frac{1}{6} - \frac{1}{4} = -\frac{1}{12} < 0$ implies $G_{\text{ind}} < 0$.
- Gauge vector (per generator): $A_1^{(\text{eff})} = -\frac{2}{3} < 0$ implies $G_{\text{ind}} < 0$ for that sector alone.

8.2 Python (pedagogical)

```
# Python: compute G_ind (units 1/Lambda^2) under the appendix convention
import numpy as np
```

```
def A1_scalar(xi=0.0):
    return 1.0/6.0 - xi
```

```

def A1_dirac():
    return 1.0/3.0

def A1_vector_plus_ghosts_per_generator():
    return -2.0/3.0

def G_ind_from_A1(A1_total, Lambda):
    if abs(A1_total) < 1e-12:
        return np.inf
    return (2.0*np.pi / A1_total) / (Lambda**2)

Lambda = 1.0e3
cases = [
    ("scalar_minimal (xi=0)", A1_scalar(0.0)),
    ("scalar_xi_1/4", A1_scalar(0.25)),
    ("1 Dirac only", A1_dirac()),
    ("1 vector+ghosts (per gen.)", A1_vector_plus_ghosts_per_generator()),
    ("scalar + Dirac", A1_scalar(0.0) + A1_dirac()),
    ("scalar + vector+ghosts", A1_scalar(0.0) + A1_vector_plus_ghosts_per_generator()),
]

print(f"{'CASE':<30} | {'A1_total':>9} | {'G_ind (1/L^2)':>14} | SIGN")
print("-"*74)
for name, A1t in cases:
    G = G_ind_from_A1(A1t, Lambda)
    sign = "ATR (G>0)" if G > 0 else ("REP (G<0)" if G != np.inf else "indet")
    print(f"{'name':<30} | {'A1t':9.4f} | {'G':14.4e} | {sign}")

```

9 Regularization note

The term proportional to Λ^2 is regulator-dependent: it appears explicitly with a Schwinger cutoff, while in dimensional regularization power divergences typically do not appear as poles. What is robust here is the local tensorial structure (controlled by a_1) and relative normalizations between species within a fixed scheme. The physical Newton constant G_{phys} requires a counterterm and a renormalization condition.

References

- [1] L. Eriksson, *Geometry, Membranes, and Life as a Resource Boundary: A Correct-by-Construction Operational Pipeline from Static Suppression to Maintenance Costs*, ai.viXra:2512.0085 (2025), <https://ai.vixra.org/abs/2512.0085>.
- [2] L. Eriksson, *The Maintenance Constraint: How Resource Boundaries Shape Cognitive Availability*, ai.viXra:2512.0084 (2025), <https://ai.vixra.org/abs/2512.0084>.
- [3] L. Eriksson, *Operational Coherence Maintenance and the Quantum–Classical Boundary: Formal Definitions, Falsifiable Protocols, and an Outlook for Cognitive Systems*, ai.viXra:2512.0081 (2025), <https://ai.vixra.org/abs/2512.0081>.
- [4] L. Eriksson, *Finite-Dimensional Davies Interface Lemmas and TFIM Witness Tests for the Heisenberg Cut as a Resource Boundary*, ai.viXra:2512.0073 (2025), <https://ai.vixra.org/abs/2512.0073>.
- [5] L. Eriksson, *The Rate Inheritance Principle: From Static Correlations to Dynamical Decoherence Rates*, ai.viXra:2512.0072 (2025), <https://ai.vixra.org/abs/2512.0072>.
- [6] L. Eriksson, *Operational Coherence Maintenance: Proven Results, Conditional Interfaces, and Open Dynamical Gaps*, ai.viXra:2512.0071 (2025), <https://ai.vixra.org/abs/2512.0071>.
- [7] L. Eriksson, *Stress Testing the Rate Inheritance Principle: Spectral Decoherence Rates and an Operational Resource Horizon*, ai.viXra:2512.0070 (2025), <https://ai.vixra.org/abs/2512.0070>.
- [8] L. Eriksson, *The Heisenberg Cut as a Resource Boundary: An Operational Outlook from Coherence Maintenance Costs*, ai.viXra:2512.0064 (2025), <https://ai.vixra.org/abs/2512.0064>.
- [9] L. Eriksson, *The Conditional Maintenance Work Theorem: Operational Power Lower Bounds from Energy Pinching and a Split-Inclusion Blueprint for Type III AQFT*, ai.viXra:2512.0061 (2025), <https://ai.vixra.org/abs/2512.0061>.
- [10] L. Eriksson, *Clustering, Recovery, and Locality in Algebraic Quantum Field Theory: Quantitative Bounds via Split Inclusions and Modular Theory*, ai.viXra:2512.0060 (2025), <https://ai.vixra.org/abs/2512.0060>.
- [11] P. B. Gilkey, *Invariance Theory, the Heat Equation, and the Atiyah–Singer Index Theorem*, CRC Press (1995).
- [12] D. V. Vassilevich, Heat kernel expansion: user’s manual, *Phys. Rept.* 388 (2003) 279–360, <https://arxiv.org/abs/hep-th/0306138>.
- [13] I. G. Avramidi, *Heat Kernel and Quantum Gravity*, Lecture Notes in Physics (2000).
- [14] A. D. Sakharov, Vacuum quantum fluctuations in curved space and the theory of gravitation, *Dokl. Akad. Nauk SSSR* 177 (1967) 70–71.
- [15] A. O. Barvinsky and G. A. Vilkovisky, The generalized Schwinger–DeWitt technique in gauge theories and quantum gravity, *Phys. Rept.* 119 (1985) 1–74.