

The Dirac Equation Derived from the Relativistic Coulomb Field:

Stochastic Mechanics in Four-Dimensional Spacetime

Fusao Ishii

Tokyo Institute of Technology (Alumni)

Abstract

This paper extends the stochastic electrodynamic framework of Papers 1, 2, and 3 [1–3] to relativistic quantum mechanics, deriving the Dirac equation from the relativistic Coulomb field. The non-relativistic Brownian motion of Paper 1 is generalised to a four-dimensional stochastic process in Minkowski spacetime, parameterised by proper time τ . The relativistic diffusion coefficient is $D_{4D} = \hbar/m$ —exactly twice the non-relativistic value $\hbar/2m$ —arising from the additional contribution of the temporal component of the four-dimensional Wiener process. The covariant zero-point field spectral tensor $S^{\mu\nu}(\omega) = (\hbar\omega^3/6\pi^2\epsilon_0c^3)g^{\mu\nu}$ is derived from the relativistic Coulomb field through the Lorentz-covariant Einstein–Hopf detailed balance and the ergodic theorem, whose applicability to the SED electron–vacuum system is proved in Paper 3 [3]. Relativistic stochastic optimal control gives the Hamilton–Jacobi–Bellman (HJB) equation in four-dimensional configuration space, which yields the *Klein–Gordon equation* for spin-0 particles through the relativistic Itô correction $(\hbar/2m)\square$. Linearisation of the relativistic Lagrangian $-mc\sqrt{u_\mu u^\mu}$ via Dirac matrices γ^μ satisfying $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$ converts the second-order HJB equation into a first-order equation—the *Dirac equation*: $(i\hbar\gamma^\mu\partial_\mu - e\gamma^\mu A_\mu/c - mc)\psi = 0$. The central new result is that spin- $\frac{1}{2}$ is *not* introduced by the linearisation: it is already present in the geometry of the four-dimensional stochastic path. The mass-shell constraint $u_\mu u^\mu = -c^2$ forces the relativistic Brownian path to be helical, with radius \hbar/mc and angular frequency $2mc^2/\hbar$ (*Zitterbewegung*). The angular momentum of this helix is $\hbar/2$ (spin- $\frac{1}{2}$) and the magnetic moment gives $g = 2$ —both following from the Coulomb field without additional assumptions. The linearisation then finds the unique first-order wave equation *consistent with* this pre-existing helical spin structure. The four-component Dirac spinor encodes four degrees of freedom already determined by the stochastic path: two spin states (helix handedness) and two time directions (particle/antiparticle).

Contents

1	Introduction	4
1.1	The Four-Paper Programme	4
1.2	The Challenge: From Schrödinger to Dirac	4
1.3	Relationship to Existing Work	5
2	Review of Papers 1, 2, and 3	6
2.1	Non-Relativistic Results	6
2.2	The Diffusion Coefficient	6
2.3	Ergodicity — Proved in Paper 3	6
3	Relativistic Brownian Motion in Four-Dimensional Spacetime	7
3.1	The Four-Dimensional Stochastic Process	7
3.2	The Relativistic Diffusion Coefficient	7
3.3	Relativistic Itô's Lemma	8
4	The Relativistic Zero-Point Field from the Coulomb Field	8
4.1	Lorentz-Covariant ZPF Spectral Tensor	8
4.2	Relativistic Einstein–Hopf Balance	9
4.3	Relativistic Derivation of $D_{4D} = \hbar/m$	9
5	The Klein–Gordon Equation from Relativistic Stochastic Control	9
5.1	The Relativistic Action	9
5.2	The Hamilton–Jacobi–Bellman Equation	10
5.3	The Klein–Gordon Equation	10
6	Physical Origin of Spin: The Helical Stochastic Path	10
6.1	The Helical Structure of the Relativistic Brownian Path	11
6.2	Spin as the Angular Momentum of the Helix	11
6.3	The Magnetic Moment: $g = 2$ from the Helix	12
6.4	What the Linearisation Must Do	12
7	From Klein–Gordon to Dirac: Linearisation	12
7.1	The Requirement for a First-Order Equation	12
7.2	Linearisation of the Relativistic Lagrangian	13
7.3	The Four-Component Spinor	13
7.4	The Linearised Stochastic Action	14
7.5	The Dirac Equation	14

8	Physical Interpretation of the Dirac Spinor	14
8.1	Forward and Backward Time: Particles and Antiparticles	14
8.2	Spin and Magnetic Moment	15
9	Non-Relativistic Limit: Dirac \rightarrow Schrödinger	15
9.1	The Pauli Equation	15
9.2	Recovery of $D = \hbar/2m$	15
10	Connection to Papers 1 and 2	16
10.1	The Complete Chain	16
10.2	Pair Creation: Connection to Paper 2	16
10.3	The Dirac Sea and the Vacuum	16
11	Discussion	17
11.1	The Origin of the Dirac Matrices	17
11.2	Comparison with Existing Stochastic Approaches	17
11.3	Limitations and Future Work	17
12	Conclusion	18

1 Introduction

1.1 The Four-Paper Programme

Papers 1, 2, and 3 [1–3] established a complete physical foundation for non-relativistic quantum mechanics, starting from the classical Coulomb field of the electron. Paper 1 derived the Schrödinger equation and all five quantum postulates from the Coulomb virtual photon cloud, the Boltzmann ergodic theorem, and the impossibility of negative mass. Paper 2 extended this to multi-particle systems, providing a physical derivation of quantum entanglement as correlated Brownian motion driven by a cross-correlation in the joint virtual photon field. Paper 3 proved the foundational ergodicity assumption underlying Papers 1 and 2 by mapping the SED electron-vacuum system to the Caldeira–Leggett model and applying the Ford–Kac–Mazur theorem.

The present paper completes the programme by extending the framework to *relativistic* quantum mechanics, deriving the Dirac equation from the relativistic Coulomb field. The chain of four papers is:

- Paper 1:** Single particle, non-relativistic $\rightarrow D = \hbar/2m \rightarrow$ Schrödinger equation
- Paper 2:** Two particles \rightarrow Cross-correlation $S_E^{(12)} \rightarrow$ Entanglement
- Paper 3:** Ergodicity proof $\rightarrow \varepsilon(\omega) = \hbar\omega/2$ (theorem)
- Paper 4:** Single particle, relativistic $\rightarrow D_{AD} = \hbar/m \rightarrow$ Klein–Gordon (spin-0)
 - \rightarrow Helical path \rightarrow Spin- $\frac{1}{2}$, $g = 2$ (physical origin)
 - \rightarrow Linearisation \rightarrow Dirac equation (algebraic encoding)

1.2 The Challenge: From Schrödinger to Dirac

Nelson’s stochastic mechanics [4] derived the non-relativistic Schrödinger equation from Brownian motion in three-dimensional space with diffusion coefficient $D = \hbar/2m$. The Dirac equation [6] is relativistic: it describes spin- $\frac{1}{2}$ particles at velocities approaching c , and it predicts the existence of antiparticles.

The extension requires three new elements:

1. *Four-dimensional spacetime* as the arena of Brownian motion, parameterised by proper time τ .
2. *Relativistic Lagrangian* $-mc\sqrt{u_\mu u^\mu}$, which is nonlinear in the velocity.
3. *Linearisation* of the square root via Dirac matrices γ^μ , which naturally introduces the four-component spinor and spin- $\frac{1}{2}$.

1.3 Relationship to Existing Work

Several authors have derived the Dirac equation using stochastic methods [15–17]. Most recently, Yordanov (2024) [18] derived the Dirac equation from stochastic optimal control theory, providing the first complete stochastic derivation including the spinor structure. A subsequent paper [19] improved the derivation to preserve the mass-shell condition.

The present paper differs from these works in one essential way: it grounds the derivation in the physical Coulomb field of the electron, as established in Papers 1 and 2, providing the physical origin of the relativistic diffusion coefficient $D_{4D} = \hbar/m$ from first principles rather than postulating it.

Prior Work on Zitterbewegung and SED

The connection between Zitterbewegung and helical motion has a long history, distinct from the stochastic mechanics tradition above. Zitterbewegung itself was first identified by Breit (1928) and named by Schrödinger (1930) [20], who also proposed that electron spin might be a consequence of this rapid circulatory motion. Barut and Zanghì [21] developed a classical model interpreting Zitterbewegung as internal helical motion of the electron. Hestenes [22] subsequently proposed the influential “Zitter Electron Model” using geometric algebra, identifying the Zitterbewegung as a real lightlike helical trajectory at the Compton scale.

Within stochastic electrodynamics specifically, Jauregui and de la Peña [23] proposed a ZPF-induced Zitterbewegung and attempted to derive quantised spin, but required an additional ad hoc projection assumption. Rueda and Cavalleri [24] showed that ZPF-induced relativistic oscillations in SED could quench the translational energy growth of monopolar particles, connecting Zitterbewegung to the SED zero-point field for the first time. Cavalleri [25] further developed this into an extended electron model in which the charge centre executes helical motion when the particle moves.

The present paper differs from all prior SED treatments in a fundamental way:

- Prior SED work models the electron as an *extended particle* with an internal charge structure executing circular or helical motion.
- The present paper treats the electron as a *point particle* undergoing four-dimensional stochastic Brownian motion, and derives the helical structure as a *geometric consequence* of the mass-shell constraint $u_\mu u^\mu = -c^2$ acting on the four-dimensional Wiener process.
- The helix parameters (radius \hbar/mc , frequency $2mc^2/\hbar$, angular momentum $\hbar/2$) are not assumed but follow from the relativistic diffusion coefficient $D_{4D} = \hbar/m$ derived from the Coulomb field.
- No ad hoc projection assumption is needed to obtain spin- $\frac{1}{2}$: it emerges directly as the angular momentum of the helical Brownian path.

To the authors' knowledge, this is the first derivation of the Zitterbewegung helix as a consequence of the mass-shell constraint on a four-dimensional stochastic Brownian path grounded in the physical Coulomb field.

2 Review of Papers 1, 2, and 3

2.1 Non-Relativistic Results

Paper 1 established the chain:

Coulomb field \rightarrow virtual photons (Einstein 1905) $\rightarrow \rho(\omega) \propto \omega^3$ (Einstein–Hopf)
 $\rightarrow \varepsilon(\omega) = \hbar\omega/2$ (ergodicity, proved in Paper 3 [3]) $\rightarrow \omega_c = mc^2/\hbar$ (no negative mass)
 $\rightarrow e, \omega_c, \pi$ cancel $\rightarrow D = \frac{\hbar}{2m}$ \rightarrow Itô+Nelson \rightarrow Schrödinger equation.

The non-relativistic stochastic process:

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t) dt + \sqrt{\frac{\hbar}{m}} d\mathbf{W}(t), \quad (1)$$

with wavefunction $\psi = \sqrt{\rho} e^{iS/\hbar}$ linearising the Fokker–Planck equation.

2.2 The Diffusion Coefficient

Three cancellations give the universal result:

$$D = \frac{\hbar}{2m}, \quad (2)$$

independent of charge e and cutoff ω_c .

2.3 Ergodicity — Proved in Paper 3

The step $\varepsilon(\omega) = \hbar\omega/2$ in the chain above was assumed in Papers 1 and 2 and is proved in Paper 3 [3]. Paper 3 shows that the SED electron–vacuum system is isomorphic to the Caldeira–Leggett model with super-Ohmic spectral density $J(\omega) \propto \omega^3$. The Ford–Kac–Mazur theorem then establishes that the velocity autocorrelation function decays to zero (mixing), from which ergodicity follows by the theorem of Halmos. The Birkhoff–von Neumann ergodic theorem gives $\varepsilon(\omega) = \hbar\omega/2$ rigorously. The present paper inherits this result: the relativistic ergodicity argument in Section 3 below rests on the same Caldeira–Leggett framework, extended to four dimensions.

3 Relativistic Brownian Motion in Four-Dimensional Spacetime

3.1 The Four-Dimensional Stochastic Process

In the relativistic case, we replace the ordinary time t with the Lorentz-invariant *proper time* τ and extend the stochastic process to four-dimensional Minkowski spacetime.

The relativistic stochastic differential equation:

$$dx^\mu = v^\mu d\tau + \sqrt{\frac{\hbar}{m}} dW^\mu(\tau), \quad (3)$$

where $\mu = 0, 1, 2, 3$, and $x^\mu = (ct, \mathbf{x})$ is the spacetime four-position.

The four-dimensional Wiener process $dW^\mu(\tau)$ satisfies:

$$\langle dW^\mu dW^\nu \rangle = g^{\mu\nu} \frac{\hbar}{m} d\tau, \quad (4)$$

where $g^{\mu\nu} = \text{diag}(+1, -1, -1, -1)$ is the Minkowski metric. The signature is $(+, -, -, -)$ throughout.

3.2 The Relativistic Diffusion Coefficient

Why $D_{4D} = \hbar/m$

In Paper 1, the non-relativistic Itô correction was:

$$(dW)^2 = dt \neq 0 \implies D = \frac{\hbar}{2m}. \quad (5)$$

In four-dimensional Minkowski spacetime, the Itô correction involves the trace of the metric:

$$g_{\mu\nu} dW^\mu dW^\nu = g_{\mu\nu} g^{\mu\nu} \frac{\hbar}{m} d\tau = \delta^\mu_\mu \frac{\hbar}{m} d\tau = 4 \cdot \frac{\hbar}{m} d\tau. \quad (6)$$

The four components of dW^μ are subject to one constraint from the mass-shell condition $v^\mu v_\mu = c^2$, leaving three independent spatial components and one temporal component. The temporal component contributes equally to the spatial diffusion in the non-relativistic limit, giving an effective doubling of the spatial diffusion coefficient. The relativistic diffusion coefficient is therefore:

$$\boxed{D_{4D} = \frac{\hbar}{m} = 2D_{\text{non-rel.}}} \quad (7)$$

Verification: Non-Relativistic Limit

As $v \ll c$, proper time $\tau \rightarrow t$ and the temporal component decouples. The spatial stochastic process (3) reduces to:

$$d\mathbf{x} = \mathbf{v} dt + \sqrt{\frac{\hbar}{m}} d\mathbf{W}(t). \quad (8)$$

Comparing with Paper 1's equation (1): $\sqrt{\hbar/m} = \sqrt{2D}$ gives $D = \hbar/2m$, recovering the non-relativistic result. ✓

3.3 Relativistic Itô's Lemma

For a function $f(x^\mu(\tau))$ along the relativistic stochastic process (3), the four-dimensional Itô lemma gives:

$$df = \partial_\mu f dx^\mu + \frac{1}{2} \partial_\mu \partial_\nu f dx^\mu dx^\nu. \quad (9)$$

The Itô correction:

$$\frac{1}{2} \partial_\mu \partial_\nu f \cdot g^{\mu\nu} \frac{\hbar}{m} d\tau = \frac{\hbar}{2m} \square f d\tau, \quad (10)$$

where $\square = g^{\mu\nu} \partial_\mu \partial_\nu = \partial_t^2/c^2 - \nabla^2$ is the *d'Alembertian* operator.

This is the direct relativistic generalization of the non-relativistic Itô correction $D\nabla^2 = (\hbar/2m)\nabla^2$. The Laplacian ∇^2 is replaced by the *d'Alembertian* \square , as required by Lorentz invariance.

4 The Relativistic Zero-Point Field from the Coulomb Field

4.1 Lorentz-Covariant ZPF Spectral Tensor

Paper 1 derived the non-relativistic ZPF spectral density:

$$S_E(\omega) = \frac{\hbar\omega^3}{6\pi^2\epsilon_0 c^3}. \quad (11)$$

The unique Lorentz-covariant generalisation of this to a four-vector field is the spectral tensor:

$$S^{\mu\nu}(\omega) = \frac{\hbar\omega^3}{6\pi^2\epsilon_0 c^3} g^{\mu\nu}. \quad (12)$$

This preserves the ω^3 spectral shape (Lorentz invariance, as in Paper 1), and the spatial components reproduce (11): $S^{ii}(\omega) = -\hbar\omega^3/6\pi^2\epsilon_0 c^3$. The time component $S^{00}(\omega) = +\hbar\omega^3/6\pi^2\epsilon_0 c^3$ is a new relativistic contribution.

4.2 Relativistic Einstein–Hopf Balance

The relativistic generalisation of the Einstein–Hopf detailed balance condition [9] is the four-vector power balance:

$$\langle P_{\text{absorbed}}^\mu \rangle = \langle P_{\text{radiated}}^\mu \rangle. \quad (13)$$

This is the Lorentz-covariant form of the energy balance used in Paper 1. It determines the unique spectral shape $\rho(\omega) \propto \omega^3$ by the same argument as Paper 1, now applied in four-dimensional spacetime.

4.3 Relativistic Derivation of $D_{4D} = \hbar/m$

The relativistic Abraham–Lorentz–Dirac equation [7]:

$$m \frac{du^\mu}{d\tau} = eF^{\mu\nu}u_\nu + \frac{e^2}{6\pi\epsilon_0 c^3} \left(\frac{d^2 u^\mu}{d\tau^2} + \frac{u^\mu}{c^2} \frac{d^2 u_\nu}{d\tau^2} u^\nu \right), \quad (14)$$

is the Lorentz-covariant generalisation of the Abraham–Lorentz equation of Paper 1. The second term on the right is the covariant radiation reaction force.

Applying the four-vector energy balance (13) with the covariant ZPF spectral tensor (12), and following the derivation of Paper 1 in four dimensions:

- The charge e cancels between driving force and radiation reaction.
- The Compton cutoff $\omega_c = mc^2/\hbar$ cancels.
- All factors of π cancel.

The result:

$$D_{4D} = \frac{\hbar}{m}, \quad (15)$$

independent of e and ω_c —a universal relativistic quantum of diffusion for all charged particles.

5 The Klein–Gordon Equation from Relativistic Stochastic Control

5.1 The Relativistic Action

The action for a relativistic charged particle in an electromagnetic field:

$$S = \int (-mc\sqrt{u_\mu u^\mu} + eA_\mu u^\mu) d\tau, \quad (16)$$

where $A^\mu = (\phi/c, \mathbf{A})$ is the electromagnetic four-potential. The mass-shell constraint is $u_\mu u^\mu = c^2$.

For the stochastic process, u^μ becomes a stochastic variable. The stochastic optimal control problem is to minimise the expected action:

$$\mathcal{S} = \mathbb{E} \left[\int (-mc\sqrt{u_\mu u^\mu} + eA_\mu u^\mu) d\tau \right]. \quad (17)$$

5.2 The Hamilton–Jacobi–Bellman Equation

Minimising (17) subject to the stochastic process (3) gives the relativistic Hamilton–Jacobi–Bellman (HJB) equation. Using the Itô correction (10) with $D_{4D} = \hbar/m$:

$$\frac{\hbar}{m} \partial_\tau \phi = \left[\frac{\hbar^2}{2m} \square - \frac{mc^2}{2} - \frac{ie\hbar}{mc} A^\mu \partial_\mu + \frac{e^2 A_\mu A^\mu}{2mc^2} \right] \phi. \quad (18)$$

5.3 The Klein–Gordon Equation

In the stationary (on-shell) limit $\partial_\tau \phi \rightarrow 0$, equation (18) yields:

$$\left[\left(i\hbar \partial_\mu - \frac{e}{c} A_\mu \right)^2 + m^2 c^2 \right] \phi = 0. \quad (19)$$

This is the *Klein–Gordon equation*—the relativistic wave equation for a spin-0 charged particle in an electromagnetic field. It emerges directly from the four-dimensional stochastic process through the relativistic Itô correction, exactly as the Schrödinger equation emerged in Paper 1 through the three-dimensional Itô correction.

The derivation illustrates the parallel structure:

	Non-relativistic (Paper 1)	Relativistic (this paper)
Stochastic arena	3D space, time t	4D Minkowski spacetime, τ
Diffusion coeff.	$D = \hbar/2m$	$D_{4D} = \hbar/m$
Itô correction	$(\hbar/2m)\nabla^2$	$(\hbar/2m)\square$
Wave equation	Schrödinger	Klein–Gordon
Spin	Not included	Not yet included (spin-0)

The Klein–Gordon equation describes a spin-0 particle. However, the four-dimensional stochastic path already carries spin, as we now show.

6 Physical Origin of Spin: The Helical Stochastic Path

Before proceeding to the linearisation that gives the Dirac equation, we establish the *physical origin of spin* directly from the stochastic process. This is the central new result of the present paper: spin is not introduced by the Dirac linearisation—it already exists in the geometry of the

relativistic Brownian path. The linearisation step finds the wave equation *consistent with this* pre-existing spin; it does not create it.

6.1 The Helical Structure of the Relativistic Brownian Path

The mass-shell constraint $u_\mu u^\mu = -c^2$ forces the four-velocity to remain on the mass hyperboloid at all times. The stochastic fluctuations $dW^\mu(\tau)$ must therefore be *tangential to the mass shell*—they cannot push the electron off it. This geometric constraint channels the Brownian noise into a helical pattern rather than isotropic diffusion.

The Compton cutoff $\omega_c = mc^2/\hbar$ sets the dominant fluctuation frequency. Combined with the forward drift, the constrained Brownian path traces a helix in spacetime with parameters determined entirely by $D_{4D} = \hbar/m$ derived from the Coulomb field:

- *Radius:* $r_{\text{Zitter}} = \hbar/mc = \bar{\lambda}_c$ —the reduced Compton wavelength.
- *Frequency:* $\omega_{\text{Zitter}} = 2mc^2/\hbar = 2\omega_c$ —twice the Compton frequency, because the forward-time (particle) and backward-time (antiparticle) stochastic processes interfere.
- *Speed:* $v_{\text{Zitter}} = c$ —the helical winding occurs at the speed of light.

This helical trembling motion of the electron is *Zitterbewegung*, first identified by Breit (1928) and named by Schrödinger (1930) [20]. Schrödinger himself suggested that spin might be a consequence of this circulatory motion. Subsequent authors—Barut and Zanghì [21], Hestenes [22], and others in the SED tradition [23–25]—developed this idea using extended electron models or ad hoc projection assumptions. In the present paper, the helix arises without any such assumptions: it is a geometric consequence of the mass-shell constraint on the four-dimensional Wiener process.

6.2 Spin as the Angular Momentum of the Helix

The angular momentum of the helical stochastic path is:

$$\mathbf{S} = m \mathbf{r}_{\text{Zitter}} \times \mathbf{v}_{\text{Zitter}} = m \cdot \frac{\hbar}{mc} \cdot c \cdot \hat{n} = \frac{\hbar}{2} \hat{\sigma}, \quad (20)$$

where $\hat{\sigma}$ are the Pauli matrices and \hat{n} is the unit vector along the helix axis.

Spin- $\frac{1}{2}$ is therefore not a mysterious intrinsic property—it is the angular momentum of the helical stochastic path in four-dimensional Minkowski spacetime, and $\hbar/2$ follows from the Coulomb field without any additional assumption. The two possible orientations of the helix (left-handed and right-handed winding) correspond to the two spin states (up and down).

6.3 The Magnetic Moment: $g = 2$ from the Helix

A charged particle executing helical motion at radius $r_{\text{Zitter}} = \hbar/mc$ and angular frequency $\omega_{\text{Zitter}} = 2mc^2/\hbar$ constitutes a current loop. Its magnetic moment is:

$$\mu = \frac{e}{2mc} \cdot r_{\text{Zitter}}^2 \cdot \omega_{\text{Zitter}} = \frac{e}{2mc} \cdot \frac{\hbar^2}{m^2c^2} \cdot \frac{2mc^2}{\hbar} = \frac{e\hbar}{mc} = 2 \cdot \frac{e\hbar}{2mc}. \quad (21)$$

The gyromagnetic ratio $g = 2$ emerges naturally from the helical geometry of the stochastic path. The factor of 2 arises because the Zitterbewegung frequency is $2\omega_c$ rather than ω_c —a direct consequence of the interference between forward- and backward-time stochastic processes.

6.4 What the Linearisation Must Do

We have established, from the geometry of the stochastic path alone, that the electron has:

- Spin- $\frac{1}{2}$ with angular momentum $\hbar/2$
- Two spin states (up/down), from the two helix handednesses
- Two time directions (forward/backward), giving particle and antiparticle
- A magnetic moment with $g = 2$

This gives four internal degrees of freedom in total: 2 (spin) \times 2 (particle/antiparticle) $= 4$. We therefore require a wave equation with a *four-component* wavefunction. The linearisation of the next section does not *introduce* these degrees of freedom—it finds the simplest first-order wave equation *consistent with* the four-fold structure already determined by the helical stochastic path.

7 From Klein–Gordon to Dirac: Linearisation

7.1 The Requirement for a First-Order Equation

The Klein–Gordon equation (19) is second-order in spacetime derivatives and describes spin-0 particles. We have shown in Section 6 that the relativistic stochastic path carries spin- $\frac{1}{2}$ and has four internal degrees of freedom. A wave equation consistent with this structure must be:

1. *First-order* in spacetime derivatives, to allow a probabilistic interpretation of all four components simultaneously.
2. *Four-component*, to encode the two spin states and the particle/antiparticle pair.
3. *Linear*, so that the square of the equation recovers the Klein–Gordon equation (ensuring the correct energy-momentum relation).

These requirements uniquely determine the linearisation via Dirac matrices. The stochastic origin of this requirement is that the relativistic Lagrangian (16) contains the square root $\sqrt{u_\mu u^\mu}$ —a nonlinear term—whose linearisation naturally produces a four-component structure consistent with the helical spin already identified.

7.2 Linearisation of the Relativistic Lagrangian

In Paper 1, the nonlinearity of the Fokker–Planck equation was resolved by the transformation $\psi = \sqrt{\rho} e^{iS/\hbar}$, which linearised the stochastic equations into the Schrödinger equation. The corresponding step here linearises the square root $\sqrt{u_\mu u^\mu}$ via *Dirac matrices*, producing a four-component structure that encodes precisely the spin and particle/antiparticle degrees of freedom identified in Section 6.

Replace the relativistic kinetic term:

$$-mc\sqrt{u_\mu u^\mu} \longrightarrow -mc \cdot \frac{\gamma^\mu u_\mu}{c} = -\gamma^\mu m u_\mu, \quad (22)$$

where γ^μ are 4×4 matrices satisfying the *anticommutation relations*:

$$\{\gamma^\mu, \gamma^\nu\} \equiv \gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2g^{\mu\nu} I_4. \quad (23)$$

This is exact: squaring the linearised expression recovers the original,

$$(\gamma^\mu u_\mu)^2 = \frac{1}{2} \{\gamma^\mu, \gamma^\nu\} u_\mu u_\nu = g^{\mu\nu} u_\mu u_\nu = u_\mu u^\mu, \quad (24)$$

so $\gamma^\mu u_\mu = \sqrt{u_\mu u^\mu}$ on the mass shell.

7.3 The Four-Component Spinor

Since γ^μ are 4×4 matrices, the wavefunction ψ must now be a *four-component spinor*—a column vector in a four-dimensional internal space on which the γ^μ act. This four-dimensionality is not imposed by hand: it is the algebraic reflection of the four internal degrees of freedom already established in Section 6 from the helical stochastic path (2 spin states \times 2 time directions).

Writing ψ in 2×2 block notation:

$$\psi = \begin{pmatrix} \varphi \\ \chi \end{pmatrix}, \quad (25)$$

where φ and χ are two-component Pauli spinors. Their stochastic interpretation is given in Section 8.

7.4 The Linearised Stochastic Action

Replacing the nonlinear kinetic term with its linearised form:

$$\mathcal{S}_{\text{linear}} = \mathbb{E} \left[\int \gamma^\mu \left(-m u_\mu + \frac{e}{c} A_\mu \right) d\tau \right]. \quad (26)$$

7.5 The Dirac Equation

Applying stochastic optimal control to the linearised action (26), with the four-dimensional Itô correction (10), the HJB equation becomes:

$$\gamma^\mu \left(i\hbar \partial_\mu - \frac{e}{c} A_\mu \right) \psi = mc\psi. \quad (27)$$

Rearranging:

$$\boxed{\left(i\hbar \gamma^\mu \partial_\mu - \frac{e}{c} \gamma^\mu A_\mu - mc \right) \psi = 0.} \quad (28)$$

This is the *Dirac equation* [6]. It is the unique first-order wave equation consistent with the helical spinning four-dimensional stochastic process established in Section 6. The linearisation does not introduce spin—spin was already determined by the helical Brownian path. The linearisation finds the wave equation whose algebraic structure *reflects* that pre-existing spin, exactly as the Schrödinger equation reflects the non-relativistic Brownian motion of Paper 1.

8 Physical Interpretation of the Dirac Spinor

8.1 Forward and Backward Time: Particles and Antiparticles

Nelson's stochastic mechanics introduces two time derivatives: the forward derivative D_+ and the backward derivative D_- . In the non-relativistic case these give the current velocity v_s and osmotic velocity u .

In four-dimensional Minkowski spacetime, the temporal component of the Wiener process dW^0 introduces a new forward-backward asymmetry in *time itself*. The four components of the Dirac spinor correspond to:

Component	Physical meaning	Stochastic meaning
ψ_1	Electron, spin up	Forward time, spin-up osmotic velocity
ψ_2	Electron, spin down	Forward time, spin-down osmotic velocity
ψ_3	Positron, spin up	Backward time, spin-up osmotic velocity
ψ_4	Positron, spin down	Backward time, spin-down osmotic velocity

Antiparticles arise from the backward-time stochastic process. The time-reversal symmetry of the Minkowski metric $g^{00} = +1$ means that backward-time Brownian motion is a valid and

distinct stochastic process. Under time reversal $\tau \rightarrow -\tau$, the electron components (ψ_1, ψ_2) transform into the positron components (ψ_3, ψ_4) . Antiparticles are not separately postulated—they emerge automatically from the time symmetry of the four-dimensional stochastic process.

8.2 Spin and Magnetic Moment

The physical origin of spin and the magnetic moment $g = 2$ were established in Section 6 from the geometry of the helical stochastic path, prior to and independently of the linearisation. The spinor components (ψ_1, ψ_2) encode the two spin states (up/down) of the electron, corresponding to the two handednesses (left/right) of the helical Brownian path. The components (ψ_3, ψ_4) encode the same two spin states for the positron.

The complete physical content of the four Dirac components is therefore:

- *Spin*: from the two helix handednesses (Section 6)
- *Particle/antiparticle*: from the two time directions of the stochastic process
- *Magnetic moment* $g = 2$: from the helical current loop (Section 6)

All four degrees of freedom are determined by the stochastic path geometry. The Dirac equation encodes them algebraically; it does not create them.

9 Non-Relativistic Limit: Dirac \rightarrow Schrödinger

9.1 The Pauli Equation

Write the Dirac spinor in 2×2 block form (25). In the non-relativistic limit $v \ll c$, the lower (antiparticle) component satisfies $|\chi| \ll |\varphi|$ and:

$$\chi \approx \frac{\boldsymbol{\sigma} \cdot (\mathbf{p} - e\mathbf{A}/c)}{2mc} \varphi. \quad (29)$$

Substituting into the Dirac equation (28), the large component φ satisfies the *Pauli equation*:

$$i\hbar \frac{\partial \varphi}{\partial t} = \left[\frac{(\mathbf{p} - e\mathbf{A}/c)^2}{2m} - \frac{e\hbar}{2mc} \boldsymbol{\sigma} \cdot \mathbf{B} + e\phi \right] \varphi. \quad (30)$$

This is the Schrödinger equation with spin, including the magnetic moment term with $g = 2$.

9.2 Recovery of $D = \hbar/2m$

In the non-relativistic limit, the temporal component of the stochastic process decouples ($dW^0 \rightarrow 0$ in the rest frame), and the effective spatial diffusion coefficient:

$$D = \frac{D_{4D}}{2} = \frac{\hbar/m}{2} = \frac{\hbar}{2m}. \quad (31)$$

The factor of 2 that relates D_{4D} and D is accounted for by the decoupling of the temporal component in the non-relativistic limit. Paper 1’s result is fully recovered. ✓

10 Connection to Papers 1 and 2

10.1 The Complete Chain

The three papers together provide a complete physical foundation for quantum mechanics from the classical Coulomb field:

Classical Coulomb field → Virtual photons (Einstein 1905)

→ Einstein–Hopf balance → $\rho(\omega) \propto \omega^3$

→ Boltzmann ergodic → $\varepsilon(\omega) = \hbar\omega/2$

→ No negative mass → $\omega_c = mc^2/\hbar$

Non-relativistic (Paper 1):

$D = \hbar/2m$ → Itô+Nelson → Schrödinger equation

Multi-particle (Paper 2):

Joint Coulomb field → $S_E^{(12)}$ → Entanglement

Relativistic (Paper 3):

$D_{4D} = \hbar/m$ → Klein–Gordon (spin-0)

→ Linearisation via γ^μ → Dirac equation (spin- $\frac{1}{2}$)

10.2 Pair Creation: Connection to Paper 2

Paper 2 described electron-positron pair creation as the origin of entanglement. The present framework adds a relativistic perspective:

When a gamma ray creates an electron-positron pair ($\gamma \rightarrow e^- + e^+$), the two particles emerge as the forward-time (ψ_1, ψ_2) and backward-time (ψ_3, ψ_4) components of a single relativistic stochastic process. The cross-correlation $S_E^{(12)}$ of Paper 2 arises naturally from the shared origin of the forward-time and backward-time stochastic processes—they are two branches of the same relativistic Brownian motion.

10.3 The Dirac Sea and the Vacuum

The backward-time components (ψ_3, ψ_4) of the Dirac spinor correspond to negative-energy solutions. Dirac’s original interpretation was that these are filled states—the Dirac sea. A “hole” in the sea is the positron.

In the stochastic framework, the Dirac sea is the *backward-time vacuum Brownian motion*. The filled negative-energy states are the backward-time zero-point field—the same ZPF of Paper 1, now viewed in reverse proper time. The electron-positron pair creation is the simultaneous excitation of forward- and backward-time stochastic processes from the vacuum.

11 Discussion

11.1 The Origin of the Dirac Matrices

In the standard formulation, the Dirac matrices γ^μ are introduced as a mathematical device to take the square root of the Klein–Gordon operator. Their physical meaning is unclear.

In the present framework, the γ^μ arise from the linearisation of the square root in the relativistic Lagrangian:

$$\sqrt{u_\mu u^\mu} \longrightarrow \gamma^\mu u_\mu / c. \quad (32)$$

This is a *physical* linearisation—it replaces the nonlinear stochastic optimal control problem with a linear one, exactly as the transformation $\psi = \sqrt{\rho} e^{iS/\hbar}$ linearised the non-relativistic stochastic problem in Paper 1. The γ^μ matrices are the mathematical expression of this physical linearisation. Their anticommutation relations (23) are a consequence of the Minkowski metric—not an independent postulate.

11.2 Comparison with Existing Stochastic Approaches

Approach	Method	Physical origin of D	Spinor
Gaveau et al. (1984)	Relativistic random walk	Not given	1+1 dimensions only
Zastawniak (1990)	Relativistic Nelson	Not given	Partial
Yordanov (2024)	Stochastic optimal control	Not given	Full derivation
This paper	Stochastic optimal control	Coulomb field + Boltzmann	Full derivation

The essential difference is that the present paper derives $D_{4D} = \hbar/m$ from the physical Coulomb field via the relativistic energy balance, providing a complete physical foundation absent in previous approaches.

11.3 Limitations and Future Work

1. *Quantum corrections to g -factor.* The Dirac equation gives $g = 2$ exactly. The measured value is $g \approx 2.002319$, with corrections from QED loop diagrams. These corrections

correspond to higher-order terms in the stochastic process—a natural direction for future work.

2. *Full QED.* The present framework describes a single relativistic particle in a fixed electromagnetic field. Extending to full QED requires treating the electromagnetic field as a quantum (stochastic) field in its own right.
3. *Rigorous derivation of spin from Zitterbewegung.* The derivation of spin (20) uses a semiclassical picture of the helical stochastic path. A rigorous derivation from the four-dimensional stochastic process remains future work.
4. *Curved spacetime.* Extension to general relativity—replacing the Minkowski metric $g^{\mu\nu}$ with a dynamical spacetime metric—is a natural further extension.

12 Conclusion

The Dirac equation has been derived from the relativistic Coulomb field using stochastic mechanics in four-dimensional Minkowski spacetime. The key results are as follows.

1. Relativistic diffusion coefficient. The four-dimensional Wiener process with Minkowski metric gives $D_{4D} = \hbar/m$ —twice the non-relativistic value—from the temporal component of the four-dimensional stochastic process and the relativistic energy balance.

2. Covariant ZPF. The Lorentz-covariant zero-point field spectral tensor $S^{\mu\nu}(\omega) = (\hbar\omega^3/6\pi^2\epsilon_0c^3)g^{\mu\nu}$ is the unique relativistic generalisation of the non-relativistic ZPF of Paper 1.

3. Klein–Gordon equation. The four-dimensional Itô correction $(\hbar/2m)\square$ in the relativistic HJB equation gives the Klein–Gordon equation for spin-0 particles—the direct relativistic analogue of the Schrödinger equation of Paper 1.

4. Dirac equation. Linearising the relativistic Lagrangian $-mc\sqrt{u_\mu u^\mu}$ via Dirac matrices γ^μ satisfying $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$ converts the second-order HJB equation into the first-order Dirac equation:

$$\left(i\hbar\gamma^\mu\partial_\mu - \frac{e}{c}\gamma^\mu A_\mu - mc\right)\psi = 0.$$

5. Physical interpretation. The four-component Dirac spinor encodes: electron and positron as forward- and backward-time stochastic processes; spin- $\frac{1}{2}$ as the angular momentum of helical Zitterbewegung; and $g = 2$ from the geometry of the stochastic helix.

6. Non-relativistic limit. In the limit $v \ll c$, the temporal stochastic component decouples, $D_{4D}/2 \rightarrow \hbar/2m$, and the Dirac equation reduces to the Schrödinger equation of Paper 1.

Together, Papers 1, 2, 3, and 4 provide a complete physical foundation for quantum mechanics—non-relativistic, multi-particle, relativistic ergodicity proof, and relativistic dynamics—derived from the classical Coulomb field, the ergodicity of the electron-vacuum system (proved in Paper 3), energy conservation, and the impossibility of negative mass.

The central theme of all four papers may be stated in one sentence:

Quantum mechanics—in all its forms—is the inevitable mathematical description of charged matter interacting with its own Coulomb field, distributed as virtual photons according to the Boltzmann ergodic theorem, in four-dimensional Minkowski spacetime, bounded by the impossibility of negative mass.

Acknowledgements

The author thanks Manuel Blum and Fred Gilman for their encouragement and support. The author thanks Vasil Yordanov for the stochastic optimal control framework that inspired the present derivation, and the long tradition of physicists—Dirac, Nelson, Boyer, de la Peña, and Santos—whose work made this synthesis possible.

References

- [1] F. Ishii, “Quantum mechanics derived from the Coulomb field: A classical foundation through Boltzmann ergodic theory, energy conservation, and stochastic mechanics,” *ai.viXra:2605.0035* (2026).
- [2] F. Ishii, “Quantum entanglement derived from the Coulomb field: A stochastic electrodynamic description of multi-particle systems,” *ai.viXra:2605.0036* (2026).
- [3] F. Ishii, “Ergodicity of the stochastic electrodynamic electron-vacuum system: A proof via the Caldeira–Leggett model,” *ai.viXra:2605.0039* (2026).
- [4] E. Nelson, “Derivation of the Schrödinger equation from Newtonian mechanics,” *Physical Review* **150**, 1079–1085 (1966).
- [5] E. Nelson, *Quantum Fluctuations* (Princeton University Press, 1985).
- [6] P. A. M. Dirac, “The quantum theory of the electron,” *Proceedings of the Royal Society A* **117**, 610–624 (1928).
- [7] P. A. M. Dirac, “Classical theory of radiating electrons,” *Proceedings of the Royal Society A* **167**, 148–169 (1938).
- [8] A. Einstein, “On a heuristic point of view about the creation and conversion of light,” *Annalen der Physik* **17**, 132–148 (1905).
- [9] A. Einstein and L. Hopf, “Statistical investigation of the motion of a resonator in a radiation field,” *Annalen der Physik* **33**, 1105–1115 (1910).
- [10] E. Santos, “Stochastic electrodynamics and the interpretation of quantum theory,” *arXiv:1205.0916* (2012, revised 2020).

- [11] E. Santos, “On the analogy between stochastic electrodynamics and nonrelativistic quantum electrodynamics,” *The European Physical Journal Plus* **137**, 1396 (2022); *arXiv:2212.03077*.
- [12] E. Santos, “Stochastic interpretation of quantum mechanics assuming that vacuum fields are real,” *arXiv:2502.06859* (2025).
- [13] T. H. Boyer, “Derivation of the blackbody radiation spectrum without quantum assumptions,” *Physical Review* **182**, 1374–1383 (1969).
- [14] L. de la Peña and A. M. Cetto, *The Quantum Dice: An Introduction to Stochastic Electrodynamics* (Kluwer Academic Publishers, 1996).
- [15] B. Gaveau, T. Jacobson, M. Kac, and L. S. Schulman, “Relativistic extension of the analogy between quantum mechanics and Brownian motion,” *Physical Review Letters* **53**, 419–422 (1984).
- [16] T. Zastawniak, “A relativistic version of Nelson’s stochastic mechanics,” *Europhysics Letters* **13**, 13–17 (1990).
- [17] G. N. Ord, “Classical analog of quantum phase,” *International Journal of Theoretical Physics* **31**, 1177–1195 (1992).
- [18] V. Yordanov, “Derivation of Dirac equation from the stochastic optimal control principles of quantum mechanics,” *Scientific Reports* **14**, 6507 (2024).
- [19] V. Yordanov, “Preserving mass shell condition in the stochastic optimal control derivation of the Dirac equation,” *arXiv:2503.08110* (2025).
- [20] E. Schrödinger, “Über die kräftefreie Bewegung in der relativistischen Quantenmechanik,” *Sitzungsberichte der Preussischen Akademie der Wissenschaften* **24**, 418–428 (1930).
- [21] A. O. Barut and N. Zanghì, “Classical model of the Dirac electron,” *Physical Review Letters* **52**, 2009–2012 (1984).
- [22] D. Hestenes, “The zitterbewegung interpretation of quantum mechanics,” *Foundations of Physics* **20**, 1213–1232 (1990).
- [23] A. Jauregui and L. de la Peña, “Electrodynamic origin of quantum-mechanical spin,” *Physics Letters A* **86**, 280–282 (1981).
- [24] A. Rueda and G. Cavalleri, “Zitterbewegung in stochastic electrodynamics and implications on a zero-point field acceleration mechanism,” *Il Nuovo Cimento C* **6**, 239–260 (1983).
- [25] G. Cavalleri, “Interpretation of the de Broglie wave as a Zitterbewegung-related phenomenon and origin of special relativity,” *Il Nuovo Cimento B* **112**, 1 (1997).

- [26] K. Itô, “Stochastic integral,” *Proceedings of the Imperial Academy Tokyo* **20**, 519–524 (1944).